

Blood Alcohol Content	Predictable Effects
.02% to .04%	<ul style="list-style-type: none"> • Light-headed • Relaxation • Sensation of warmth • Altered mood ("high") • Minor impairment of judgment
.05% to .07%	<ul style="list-style-type: none"> • Buzzed • Relaxation • Euphoria • Lower inhibitions • Minor impairment of reasoning and memory • Exaggerated emotions (both good and bad)
.08% to .10%	<ul style="list-style-type: none"> • Legally impaired • Euphoria • Fatigue • Impairment in: <ul style="list-style-type: none"> ◦ Balance ◦ Speech ◦ Vision ◦ Reaction time and hearing • Judgment and self-control are impaired
.11% to .15%	<ul style="list-style-type: none"> • Drunk • "High" reduced • Depressive effect more pronounced (anxiety, depression or unease) • Gross motor impairment • Judgment and perception severely impaired
.16% to .19%	<ul style="list-style-type: none"> • Very drunk • Strong state of depression • Nausea • Disorientation • Dizzy • Increased motor impairment • Blurred vision • Judgment further impaired
.20% to .24%	<ul style="list-style-type: none"> • Dazed and confused • Gross disorientation to time and place • Increased nausea and vomiting • May need assistance to stand/walk • Impervious to pain

	<ul style="list-style-type: none"> ● Blackout likely
.25% to .30%	<ul style="list-style-type: none"> ● Stupor ● All mental, physical and sensory functions are severely impaired ● Accidents very likely ● Little comprehension ● May pass out suddenly
.31% and up	<ul style="list-style-type: none"> ● Coma ● Level of surgical amnesia ● Onset of coma ● Possibility of acute alcohol poisoning ● Death due to respiratory arrest is likely in 50% of drinkers